arXiv:1210.8140v2 [math.SG] 1 Nov 2012

A NOTE ON THE FRONT SPINNING CONSTRUCTION

ROMAN GOLOVKO

ABSTRACT. In this paper we introduce a notion of front S^m-spinning for Legendrian submanifolds of $\mathbb{R}^{2n+1}$. It generalizes the notion of front S^1-spinning which was invented by Ekholm, Etnyre and Sullivan in [9]. We use it to prove that there are infinitely many pairs of exact Lagrangian cobordant and not pairwise Legendrian isotopic Legendrian $S^1 \times S^{i_1} \times \cdots \times S^{i_k}$ in $\mathbb{R}^{2(\sum_{j=1}^{k} i_j+1)+1}$ which have the same classical invariants if one of i_j's is odd.

1. INTRODUCTION AND MAIN RESULTS

Basic definitions. *Standard contact* $(2n+1)$*-dimensional space* is Euclidean space $\mathbb{R}^{2n+1}$ equipped with the completely non-integrable field of hyperplanes $\xi = \ker \alpha$, where α is the contact 1-form $\alpha = dz - \sum_{i=1}^{n} y_i dx_i$ in Euclidean coordinates $(x_1, y_1, \dots, x_n, y_n, z)$. The *Reeb vector field* R_α which corresponds to $\alpha = dz - \sum_{i=1}^{n} y_i dx_i$ is given by $R_\alpha = \frac{\partial}{\partial z}$. An immersion of an n-manifold into $\mathbb{R}^{2n+1}$ is *Legendrian* if it is everywhere tangent to the hyperplane field ξ, and the image of a Legendrian embedding is a *Legendrian submanifold*. The *Reeb chords* of a Legendrian submanifold Λ are segments of flow lines of R_α starting and ending at points of Λ. The *symplectization* of $\mathbb{R}^{2n+1}$ is the exact symplectic manifold $(\mathbb{R} \times \mathbb{R}^{2n+1}, d(e^t\alpha))$, where t is a coordinate on $\mathbb{R}$. There are two natural projections

$$\Pi_F(x_1, y_1, \dots, x_n, y_n, z) = (x_1, \dots, x_n, z) \quad \text{and}$$
$$\Pi_L(x_1, y_1, \dots, x_n, y_n, z) = (x_1, y_1, \dots, x_n, y_n)$$

that we call the *front projection* and the *Lagrangian projections*, respectively. The Lagrangian projection $\Pi_L(\Lambda)$ of a Legendrian submanifold Λ is an exact Lagrangian immersion into $\mathbb{R}^{2n}$. Note that in the generic situation, i.e. for Λ in an open dense subset of all Legendrian submanifolds with C^∞ topology, the self-intersection of $\Pi_L(\Lambda)$ consists of a finite number of transverse double points. These points correspond to all Reeb chords of Λ. A Legendrian submanifold is called *chord generic* if it has a finite number of Reeb chords. From now on we assume that all Legendrian submanifolds of $\mathbb{R}^{2n+1}$ are connected and chord generic.

Classical and non-classical invariants of Legendrian submanifolds. There are two classical invariants of a closed, orientable Legendrian submanifold $\Lambda \subset \mathbb{R}^{2n+1}$, namely the *Thurston–Bennequin invariant* (number) and the *rotation class*.

Date: November 2, 2012.

2000 *Mathematics Subject Classification.* Primary 53D12; Secondary 53D42.

Key words and phrases. Legendrian submanifold, Lagrangian cobordism, Legendrian contact homology.

The Thurston–Bennequin invariant was originally defined by Bennequin, see [2], and independently by Thurston for Legendrian knots in $\mathbb{R}^3$, and then was generalized to higher dimensions by Tabachnikov [19]. The Thurston–Bennequin number $tb(\Lambda)$ of a closed, oriented Legendrian $\Lambda \subset \mathbb{R}^{2n+1}$ is the linking number $lk(\Lambda, \Lambda')$, where Λ' is an oriented submanifold obtained from Λ by a small shift in the direction of R_α.

The rotation class $r(\Lambda)$ was defined by Ekholm, Etnyre and Sullivan for all $n \geq 1$ in [9] and is equal to the homotopy class of $(f, df_{\mathbb{C}})$ in the space of complex fiberwise isomorphisms $T\Lambda \otimes \mathbb{C} \to \xi$, where $f : \Lambda \to \mathbb{R}^{2n+1}$ is an embedding of Λ. Note that if $\Lambda = S^n$ and n is odd, then $r(\Lambda) \in \pi_n(U(n)) \simeq \mathbb{Z}$ and we call $r(\Lambda)$ the *rotation number*.

Legendrian contact homology is a non-classical invariant of a closed, orientable Legendrian submanifold of $\mathbb{R}^{2n+1}$. It was constructed in [5] for the case when $n = 1$ and in [10] for all $n \geq 1$. The Legendrian contact homology of a closed, orientable Legendrian submanifold Λ with the finite set of Reeb chords $\mathcal{C}_\Lambda$ is the homology of the noncommutative differential graded algebra $(\mathcal{A}_\Lambda, \partial_\Lambda)$ over $\mathbb{Z}_2$ freely generated by the elements of $\mathcal{C}_\Lambda$ and is denoted by $LCH_*(\Lambda)$. The differential ∂_Λ counts holomorphic curves in the symplectisation of $\mathbb{R}^{2n+1}$ whose domains are disks with points removed on the boundary. At these points, the holomorphic curve has one positive asymptotic and several negative asymptotics. For more details we refer to [10].

Note that $(\mathcal{A}_\Lambda, \partial_\Lambda)$ and even its homology may be infinite dimensional and hence it is difficult to use it for practical applications. One of the ways to extract useful information from $(\mathcal{A}_\Lambda, \partial_\Lambda)$ is to follow Chekanov's method of linearization. An augmentation ε is an algebra homomorphism from $(\mathcal{A}_\Lambda, \partial_\Lambda)$ to $(\mathbb{Z}_2, 0)$ which satisfies $\varepsilon(1) = 1$ and $\varepsilon \circ \partial_\Lambda = 0$ and allows us to linearize the differential graded algebra to a finite dimensional complex $LC^\varepsilon := (A_\Lambda, \partial^\varepsilon_{1,\Lambda})$ with homology groups $LCH^\varepsilon_*(\Lambda)$. Here A_Λ is the vector space over $\mathbb{Z}_2$ generated by the elements of $\mathcal{C}_\Lambda$. We let $LCH^*_\varepsilon(\Lambda)$ be the homology of the dual complex $LC_\varepsilon(\Lambda) := Hom(LC^\varepsilon(\Lambda), \mathbb{Z}_2)$. The linearized homology (cohomology) groups may depend on the choice of ε. However, the set of graded groups $\{LCH^\varepsilon_*(\Lambda)\}$ ($\{LCH^*_\varepsilon(\Lambda)\}$), where ε is any augmentation of $(\mathcal{A}_\Lambda, \partial_\Lambda)$, provides a Legendrian isotopy invariant, see [5].

In [9], Ekholm, Etnyre and Sullivan used Legendrian contact homology to prove that for any $n > 1$ there is an infinite family of Legendrian embeddings of the n-sphere into $\mathbb{R}^{2n+1}$ that are not Legendrian isotopic even though they have the same classical invariants. They also prove similar results for Legendrian surfaces and n-tori, see [9]. These results indicate that the theory of Legendrian submanifolds of standard contact $(2n+1)$-space is very rich.

Main results. Observe that the family of n-tori from [9] is constructed using the front S^1-spinning, which is a procedure to produce a closed, orientable Legendrian submanifold $\Sigma_{S^1}\Lambda \subset \mathbb{R}^{2n+3}$ from a closed, orientable Legendrian submanifold $\Lambda \subset \mathbb{R}^{2n+1}$. In Section 2 we introduce a notion of front S^m-spinning. It generalizes the S^1-spinning construction and produces a closed, orientable Legendrian submanifold $\Sigma_{S^m}\Lambda \subset \mathbb{R}^{2(n+m)+1}$ from a closed, orientable Legendrian submanifold $\Lambda \subset \mathbb{R}^{2n+1}$. We have to mention that the special case of the "higher dimensional spinning" construction has already appeared in the work of Ekholm, Etnyre and Sabloff, see [8]. In Section 3 we investigate the behavior of the front S^m-spinning under the relation of an embedded Lagrangian cobordism.

Observe that a closed, orientable Legendrian submanifold $\Lambda \subset \mathbb{R}^{2n+1}$ gives rise to an exact Lagrangian cylinder $C(\Lambda) = \mathbb{R} \times \Lambda \subset (\mathbb{R} \times \mathbb{R}^{2n+1}, d(e^t\alpha))$.

Definition 1.1. A Lagrangian (an exact Lagrangian) cobordism L between two closed, orientable Legendrian submanifolds $\Lambda_-, \Lambda_+ \subset \mathbb{R}^{2n+1}$ is an embedded Lagrangian (exact Lagrangian) submanifold in the symplectization of $\mathbb{R}^{2n+1}$ so that L agrees with $C(\Lambda_-)$ for $t \leq -T_L$, with $C(\Lambda_+)$ for $t \geq T_L$, $L^c := L|_{[-T_L-1, T_L+1]\times\mathbb{R}^{2n+1}}$ is compact for some $T_L \gg 0$ and we write $\Lambda_- \prec_L^{lag} \Lambda_+$ ($\Lambda_- \prec_L^{ex} \Lambda_+$). We will in general not distinguish between L and L^c and call both L. In the case when $\Lambda_- = \emptyset$, i.e. when Λ_+ has a Lagrangian (an exact Lagrangian) filling, we write $\emptyset \prec_L^{lag} \Lambda_+$ ($\emptyset \prec_L^{ex} \Lambda_+$).

From now on we assume that all Lagrangian cobordisms in the symplectization of $\mathbb{R}^{2n+1}$ are orientable.

Our goal is to prove the following result:

Proposition 1.2. *Let Λ_-, Λ_+ be two closed, orientable Legendrian submanifolds of $\mathbb{R}^{2n+1}$. If $\Lambda_- \prec_L^{lag} \Lambda_+$, then there exists a Lagrangian cobordism $\Sigma_{S^m} L$ such that $\Sigma_{S^m}\Lambda_- \prec_{\Sigma_{S^m} L}^{lag} \Sigma_{S^m}\Lambda_+$. In addition, if $\Lambda_- \prec_L^{ex} \Lambda_+$, then there exists an exact Lagrangian cobordism $\Sigma_{S^m} L$ such that $\Sigma_{S^m}\Lambda_- \prec_{\Sigma_{S^m} L}^{ex} \Sigma_{S^m}\Lambda_+$.*

We then use the fact about the relation between the linearized Legendrian contact cohomology of a Legendrian submanifold of $\mathbb{R}^{2n+1}$ and the singular homology of its exact Lagrangian filling described by Ekholm in [7] and prove the following:

Proposition 1.3. *Let Λ_- and Λ_+ be two closed, orientable Legendrian submanifolds of $\mathbb{R}^{2n+1}$ such that $\emptyset \prec_{L_{\Lambda_-}}^{ex} \Lambda_-$ and $\Lambda_- \prec_L^{ex} \Lambda_+$ with $\dim(H_i(L;\mathbb{Z}_2)) > \dim(H_i(\Lambda_-;\mathbb{Z}_2))$ for some i. Then*

1. *Λ_- is not Legendrian isotopic to Λ_+,*
2. *$\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}}\Lambda_-$ is not Legendrian isotopic to $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}}\Lambda_+$ for $i_1, \dots, i_k \geq i$, where i is the smallest number such that $\dim(H_i(L;\mathbb{Z}_2)) > \dim(H_i(\Lambda_-;\mathbb{Z}_2))$.*

In Section 6 we say a few words about the way to get a variant of Proposition 1.3 using the theory of generating families. For the basic definitions of this theory we refer to [18].

We apply Propositions 1.2 and 1.3 to a certain family of Legendrian knots and get the following theorem:

Theorem 1.4. *There are infinitely many pairs of exact Lagrangian cobordant and not pairwise Legendrian isotopic Legendrian $S^1 \times S^{i_1} \times \dots \times S^{i_k}$ in $\mathbb{R}^{2(\sum_{j=1}^{k} i_j+1)+1}$ which have the same classical invariants if one of i_j's is odd.*

2. CONSTRUCTION

In this section we define a notion of front S^m-spinning. It is a natural generalization of the front S^1-spinning invented by Ekholm, Etnyre and Sullivan in [9].

Let Λ be a closed, orientable Legendrian submanifold of $\mathbb{R}^{2n+1}$ parameterized by $f_\Lambda : \Lambda \to \mathbb{R}^{2n+1}$ with

$$f_\Lambda(p) = (x_1(p), y_1(p), \dots, x_n(p), y_n(p), z(p))$$

for $p \in \Lambda$. Without loss of generality assume that $x_1(p) > 0$ for all p. The front projection of Λ is parameterized by $\Pi_F \circ f_\Lambda$ and

$$\Pi_F \circ f_\Lambda(p) = (x_1(p), x_2(p), \dots, x_n(p), z(p)).$$

We embed $\mathbb{R}^{n+1}$ into $\mathbb{R}^{n+m+1}$ via

$$(x_1, \dots, x_n, z) \to (x_{-m+1} = 0, \dots, x_0 = 0, x_1, \dots, x_n, z)$$

and define $\Sigma_{S^m}\Lambda$ to be the Legendrian submanifold of $\mathbb{R}^{2(m+n)+1}$ whose front projection $\Pi_F(\Sigma_{S^m}\Lambda)$ is parameterized by

$$(\tilde{x}_{-m+1}(p, \theta, \overline{\phi}), \dots, \tilde{x}_1(p, \theta, \overline{\phi}), x_2(p), \dots, x_n(p), z(p)),$$

where

$$\begin{cases} \tilde{x}_{-m+1}(p, \theta, \overline{\phi}) = x_1(p) \sin\theta \sin\phi_1 \dots \sin\phi_{m-1}, \\ \tilde{x}_{-m+2}(p, \theta, \overline{\phi}) = x_1(p) \cos\theta \sin\phi_1 \dots \sin\phi_{m-1}, \\ \dots \\ \tilde{x}_1(p, \theta, \overline{\phi}) = x_1(p) \cos\phi_{m-1}, \end{cases} \tag{2.1}$$

$\theta \in [0, 2\pi)$ and $\overline{\phi} = (\phi_1, \dots, \phi_{m-1}) \in [0, \pi]^{m-1}$.

We now show the way to parametrize $\Sigma_{S^m}\Lambda$. Consider a submanifold $\tilde{\Lambda} \subset \mathbb{R}^{2(n+m)+1}$ parameterized by $f_{\tilde{\Lambda}} : \Lambda \times S^m \to \mathbb{R}^{2(n+m)+1}$ with

$$f_{\tilde{\Lambda}}(p, \theta, \overline{\phi}) = (\tilde{x}_{-m+1}(p, \theta, \overline{\phi}), \tilde{y}_{-m+1}(p, \theta, \overline{\phi}) \dots, \tilde{x}_1(p, \theta, \overline{\phi}), \tilde{y}_1(p, \theta, \overline{\phi}), x_2(p), \dots, z(p)),$$

where

$$\begin{cases} \tilde{y}_{-m+1}(p, \theta, \overline{\phi}) = y_1(p) \sin\theta \sin\phi_1 \dots \sin\phi_{m-1}, \\ \tilde{y}_{-m+2}(p, \theta, \overline{\phi}) = y_1(p) \cos\theta \sin\phi_1 \dots \sin\phi_{m-1}, \\ \dots \\ \tilde{y}_1(p, \theta, \overline{\phi}) = y_1(p) \cos\phi_{m-1}, \end{cases} \tag{2.2}$$

$\theta \in [0, 2\pi)$, $\overline{\phi} = (\phi_1, \dots, \phi_{m-1}) \in [0, \pi]^{m-1}$ and $\tilde{x}_i$'s are defined as above.

It is easy to see that

$$\begin{aligned} f^*_{\tilde{\Lambda}}(dz - \sum_{i=-m+1}^{n} y_i dx_i) &= dz(p) - \sum_{i=2}^{n} y_i(p) dx_i(p) \\ &- y_1(p)(\sin^2\phi_{m-1}(\dots(\sin^2\phi_1(\sin^2\theta + \cos^2\theta) + \cos^2\phi_1) + \dots) + \cos^2\phi_{m-1}) dx_1(p) \\ &- y_1(p)x_1(p)(\sin\theta\cos\theta - \sin\theta\cos\theta)\sin^2\phi_1 \dots \sin^2\phi_{m-1} d\theta \\ &- y_1(p)x_1(p)((\sin^2\theta + \cos^2\theta)\sin\phi_1\cos\phi_1 - \sin\phi_1\cos\phi_1)\sin^2\phi_2 \dots \sin^2\phi_{m-1} d\phi_1 \\ &- y_1(p)x_1(p)((\dots((\sin^2\theta + \cos^2\theta)\sin^2\phi_1 + \cos^2\phi_1)\dots \\ &+ \cos^2\phi_{m-2})\sin\phi_{m-1}\cos\phi_{m-1} - \sin\phi_{m-1}\cos\phi_{m-1}) d\phi_{m-1}. \end{aligned} \tag{2.3}$$

Since Λ is a Legendrian submanifold of $\mathbb{R}^{2n+1}$ and hence $f^*_\Lambda(dz - \sum_{i=1}^{n} y_i dx_i) = 0$, we use Formula 2.3 and get that

$$f^*_{\tilde{\Lambda}}(dz - \sum_{i=-m+1}^{n} y_i dx_i) = 0. \tag{2.4}$$

Since

$$f_\Lambda(p) = (x_1(p), \dots, y_n(p), z(p)),$$

where $p \in \Lambda$, is a parametrization of an embedded n-dimensional submanifold and $x_1(p) > 0$ for all $p \in \Lambda$, we easily see that

$$f_{\tilde{\Lambda}}(p, \theta, \overline{\phi}) = (\tilde{x}_{-m+1}(p, \theta, \overline{\phi}), \tilde{y}_{-m+1}(p, \theta, \overline{\phi}) \dots, \tilde{x}_1(p, \theta, \overline{\phi}), \tilde{y}_1(p, \theta, \overline{\phi}), x_2(p), \dots, z(p)) \tag{2.5}$$

with $\theta \in [0, 2\pi)$, $\overline{\phi} = (\phi_1, \dots, \phi_{m-1}) \in [0, \pi]^{m-1}$ is a parametrization of an embedded $(n+m)$-dimensional submanifold. Thus we use Formula 2.4 and get that $\tilde{\Lambda}$ is an embedded Legendrian submanifold of $\mathbb{R}^{2(n+m)+1}$ whose front projection coincides with $\Pi_F(\Sigma_{S^m}\Lambda)$. Hence $\tilde{\Lambda} = \Sigma_{S^m}\Lambda$.

3. PROOF OF PROPOSITION 1.2

Here we prove Proposition 1.2 by mimicing the proof of Proposition 1.5 from [15].

Given two closed, orientable Legendrian submanifolds $\Lambda_-, \Lambda_+ \subset \mathbb{R}^{2n+1}$ such that

$$\Lambda_\pm \subset \{(x_1, y_1, \dots, x_n, y_n, z) \in \mathbb{R}^{2n+1} \mid x_1 > 0\} \tag{3.1}$$

and $\Lambda_- \prec_L^{lag} \Lambda_+$. Let $f_L : L \to \mathbb{R}^{2n+2}$ be a parametrization of L with

$$f_L(p) = (t(p), x_1(p), y_1(p), \dots, x_n(p), y_n(p), z(p)), \tag{3.2}$$

where $p \in L$. Assume that $x_1(p) > 0$ for all p (Formula 3.1 implies that $\{f_L(p) \mid x_1(p) \leq 0\}$ is compact and hence can be translated in such a way that $x_1(p) > 0$ for all p). We now construct a Lagrangian cobordism $\Sigma_{S^m}L$ from $\Sigma_{S^m}\Lambda_-$ to $\Sigma_{S^m}\Lambda_+$. Define $\Sigma_{S^m}L$ to be parametrized by $f_{\Sigma_{S^m}L} : L \times S^m \to \mathbb{R} \times \mathbb{R}^{2(n+m)+1}$ with

$$f_{\Sigma_{S^m}L}(p, \theta, \overline{\phi}) = (t(p), \tilde{x}_{-m+1}(p, \theta, \overline{\phi}), \tilde{y}_{-m+1}(p, \theta, \overline{\phi}), \dots, \tilde{x}_1(p, \theta, \overline{\phi}), \tilde{y}_1(p, \theta, \overline{\phi}), x_2(p), \dots, z(p)),$$

where $\tilde{x}_i$'s and $\tilde{y}_i$'s are defined by Formulas 2.1 and 2.2 for x_i's and y_i's from Formula 3.2, $p \in L$, $\theta \in [0, 2\pi)$ and $\overline{\phi} = (\phi_1, \dots, \phi_{m-1}) \in [0, \pi]^{m-1}$.

Here we show that $\Sigma_{S^m}L$ is a Lagrangian cobordism from $\Sigma_{S^m}\Lambda_-$ to $\Sigma_{S^m}\Lambda_+$. We first note that $f_{\Sigma_{S^m}L}(L \times S^m) \cap \{t = t_0\}$ for $t_0 \geq T_L$ (or $t_0 \leq -T_L$) can be parametrized by $f_{\Sigma_{S^m}\Lambda_+}(p, \theta, \overline{\phi})$ (or $f_{\Sigma_{S^m}\Lambda_-}(p, \theta, \overline{\phi})$) for $f_{\Sigma_{S^m}\Lambda_\pm}(p, \theta, \overline{\phi})$ from Formula 2.5, where $p \in \Lambda_+ \subset \partial L^c$ (or $p \in \Lambda_- \subset \partial L^c$), $\theta \in [0, 2\pi)$ and $\overline{\phi} = (\phi_1, \dots, \phi_{m-1}) \in [0, \pi]^{m-1}$. In addition, from the fact that L^c is compact it follows that $\Sigma_{S^m}L^c$ is also compact. It remains to prove that the cobordism $\Sigma_{S^m}L$ is an embedded Lagrangian cobordism.

Observe that

$$
\begin{aligned}
(3.3)\qquad & f^*_{\Sigma_{S^m}L}(d(e^t(dz-\sum_{i=-m+1}^{n} y_i dx_i)))\\
&= e^t(dt(p)\wedge dz(p)-\sum_{i=2}^{n} dy_i(p)\wedge dx_i(p)-\sum_{i=2}^{n} y_i(p)dt(p)\wedge dx_i(p)\\
&-(\dots((\sin^2\theta+\cos^2\theta)\sin^2\phi_1+\cos^2\phi_1)\dots+\cos^2\phi_{m-1})dy_1(p)\wedge dx_1(p)\\
&-(\dots((\sin^2\theta+\cos^2\theta)\sin^2\phi_1+\cos^2\phi_1)\dots+\cos^2\phi_{m-1})y_1(p)dt(p)\wedge dx_1(p)\\
&\dots\\
&-((\dots((\sin^2\theta+\cos^2\theta)\sin^2\phi_1+\cos^2\phi_1)\dots+\cos^2\phi_{i-1})\sin\phi_i\cos\phi_i\\
&-\sin\phi_i\cos\phi_i)\sin^2\phi_{i+1}\dots\sin^2\phi_{m-1}y_1(p)d\phi_i\wedge dx_1(p)\\
&\dots\\
&-((\dots((\sin^2\theta+\cos^2\theta)\sin^2\phi_1+\cos^2\phi_1)\dots+\cos^2\phi_{i-1})\sin\phi_i\cos\phi_i\\
&-\sin\phi_i\cos\phi_i)\sin^2\phi_{i+1}\dots\sin^2\phi_{m-1}x_1(p)dy_1(p)\wedge d\phi_i\\
&\dots\\
&-((\dots((\sin^2\theta+\cos^2\theta)\sin^2\phi_1+\cos^2\phi_1)\dots+\cos^2\phi_{i-1})\sin\phi_i\cos\phi_i\\
&-\sin\phi_i\cos\phi_i)\sin^2\phi_{i+1}\dots\sin\phi_j\cos\phi_j\dots\sin^2\phi_{m-1}x_1(p)y_1(p)d\phi_i\wedge d\phi_j\\
&\dots\\
&-(\sin\theta\cos\theta-\sin\theta\cos\theta)\sin^2\phi_1\dots\sin^2\phi_{m-1}x_1(p)y_1(p)dt(p)\wedge d\theta\\
&\dots\\
&-(\dots((\sin^2\theta+\cos^2\theta)\sin^2\phi_1+\cos^2\phi_1)\dots)\sin^2\phi_{m-2}\\
&+\cos^2\phi_{m-2}-1)sin\phi_{m-1}\cos\phi_{m-1}y_1(p)x_1(p)dt(p)\wedge d\phi_{m-1}).
\end{aligned}
$$

In addition, since L is a Lagrangian cobordism and hence

$$
\begin{aligned}
& e^t(dt(p)\wedge dz(p)-\sum_{i=2}^{n} dy_i(p)\wedge dx_i(p)-\sum_{i=2}^{n} y_i(p)dt(p)\wedge dx_i(p))\\
&= e^t(dy_1(p)\wedge dx_1(p)+y_1(p)dt(p)\wedge dx_1(p)),
\end{aligned}
$$

we use Formula 3.3 and get that

$$
(3.4)\qquad f^*_{\Sigma_{S^m}L}(d(e^t(dz-\sum_{i=-m+1}^{n} y_i dx_i)))=0.
$$

Since

$$
f_L(p)=(t(p),x_1(p),y_1(p),\dots,x_n(p),y_n(p),z(p)),
$$

where $p \in L$, is a parametrization of an embedded $(n+1)$-dimensional cobordism and $x_1(p) > 0$ for $p \in L$, one easily sees that

$$f_{\Sigma_{S^m} L}(p,\theta,\overline{\phi}) = (t(p), \tilde{x}_{-m+1}(p,\theta,\overline{\phi}), \tilde{y}_{-m+1}(p,\theta,\overline{\phi}), \dots, \tilde{x}_1(p,\theta,\overline{\phi}), \tilde{y}_1(p,\theta,\overline{\phi}), x_2(p), \dots, z(p)),$$

where $p \in L$, $\theta \in [0, 2\pi)$ and $\overline{\phi} = (\phi_1, \dots, \phi_{m-1}) \in [0,\pi]^{m-1}$, is a parametrization of an embedded $(n+m+1)$-dimensional cobordism. Thus we use Formula 3.4 and get that $\Sigma_{S^m} L$ is really an embedded Lagrangian cobordism from $\Sigma_{S^m}\Lambda_-$ to $\Sigma_{S^m}\Lambda_+$.

We now assume that $\Lambda_- \prec_L^{ex} \Lambda_+$. Then there exists a function $h_L \in C^\infty(f_L(L), \mathbb{R})$ such that

$$dh_L = e^t(dz - \sum_{i=1}^{n} y_i dx_i).$$

We use a calculation similar to Formula 2.3 and get that

$$f^*_{\Sigma_{S^m} L}(e^t(dz - \sum_{i=-m+1}^{n} y_i dx_i)) = e^{t(p)}(dz(p) - \sum_{i=1}^{n} y_i(p) dx_i(p)). \tag{3.5}$$

Since $f_{\Sigma_{S^m} L}$ is an embedding, we can define $h_{\Sigma_{S^m} L} \in C^\infty(f_{\Sigma_{S^m} L}(\Sigma_{S^m} L), \mathbb{R})$ by setting

$$(f^*_{\Sigma_{S^m} L} h_{\Sigma_{S^m} L})(p,\theta,\overline{\phi}) := (f^*_L h_L)(p).$$

We now use Formula 3.5 and get that

$$d(f^*_{\Sigma_{S^m} L} h_{\Sigma^m L}) = e^{t(p)}(dz(p) - \sum_{i=1}^{n} y_i(p) dx_i(p)) = f^*_{\Sigma_{S^m} L}(e^t(dz - \sum_{i=-m+1}^{n} y_i dx_i)). \tag{3.6}$$

$f_{\Sigma_{S^m} L}$ is an embedding and hence Formula 3.6 implies that

$$d(h_{\Sigma_{S^m} L}) = e^t(dz - \sum_{i=-m+1}^{n} y_i dx_i).$$

Hence $\Sigma_{S^m} L$ is an exact Lagrangian cobordism. This finishes the proof of Proposition 1.2.

Note that the proof of Proposition 1.2 can be easily modified to become a proof of the following remark:

Remark 3.1. Given a closed, orientable Legendrian submanifold $\Lambda \subset \mathbb{R}^{2n+1}$. If $\emptyset \prec^{lag}_{L_\Lambda} \Lambda$ ($\emptyset \prec^{ex}_{L_\Lambda} \Lambda$), then there exists a Lagrangian (an exact Lagrangian) filling $L_{\Sigma_{S^m}\Lambda}$ such that $\emptyset \prec^{lag}_{L_{\Sigma_{S^m}\Lambda}} \Sigma_{S^m}\Lambda$ ($\emptyset \prec^{ex}_{L_{\Sigma_{S^m}\Lambda}} \Sigma_{S^m}\Lambda$).

4. PROOF OF PROPOSITION 1.3

Let Λ_- and Λ_+ be two closed, orientable Legendrian submanifolds of $\mathbb{R}^{2n+1}$ with $\emptyset \prec^{ex}_{L_{\Lambda_-}} \Lambda_-$, $\Lambda_- \prec^{ex}_L \Lambda_+$ and

$$\dim(H_i(L;\mathbb{Z}_2)) > \dim(H_i(\Lambda_-;\mathbb{Z}_2)) \tag{4.1}$$

for some i.

We first construct an exact Lagrangian filling of Λ_+. Since Λ_- is connected, and L, L_{Λ_-} are exact Lagrangian cobordisms in the symplectization of $\mathbb{R}^{2n+1}$, we glue the positive end of L_{Λ_-} to the negative end of L and get an exact Lagrangian filling L_{Λ_+} of Λ_+.

Consider the Mayer-Vietoris long exact sequence for L_{Λ_-}, L and L_{Λ_+}

$$\cdots \to H_i(\Lambda_-;\mathbb{Z}_2) \to H_i(L;\mathbb{Z}_2) \oplus H_i(L_{\Lambda_-};\mathbb{Z}_2) \xrightarrow{\Phi} H_i(L_{\Lambda_+};\mathbb{Z}_2) \to \dots . \tag{4.2}$$

From Formulas 4.1 and 4.2 it follows that

$$\begin{aligned}\dim(H_i(L_{\Lambda_+};\mathbb{Z}_2)) &\geq \dim(\Phi(H_i(L;\mathbb{Z}_2) \oplus H_i(L_{\Lambda_-};\mathbb{Z}_2)))\\ &\geq \dim(H_i(L_{\Lambda_-};\mathbb{Z}_2)) + \dim(H_i(L;\mathbb{Z}_2)) - \dim(H_i(\Lambda_-;\mathbb{Z}_2))\\ &> \dim(H_i(L_{\Lambda_-};\mathbb{Z}_2)).\end{aligned}$$

Hence we get that

$$\dim(H_i(L_{\Lambda_+};\mathbb{Z}_2)) > \dim(H_i(L_{\Lambda_-};\mathbb{Z}_2)). \tag{4.3}$$

We now define the following notations. If Λ is a closed, orientable Legendrian submanifolds of $\mathbb{R}^{2n+1}$, then we denote by $\mathcal{L}(\Lambda)$ the set of all embedded exact Lagrangian fillings of Λ and

$$\mathcal{H}^{geom}_{\Lambda} := \{(\dim(H_i(L_\Lambda;\mathbb{Z}_2)))_i : L_\Lambda \in \mathcal{L}(\Lambda)\}.$$

Here we remind the reader of the following fact described by Ekholm in [7], which comes from certain observations of Seidel in wrapped Floer homology [1], [14]:

Fact 4.1 ([7]). *Let Λ be a closed, orientable Legendrian submanifold of $\mathbb{R}^{2n+1}$ and $\emptyset \prec^{ex}_{L_\Lambda} \Lambda$. Then*

$$H_i(L_\Lambda;\mathbb{Z}_2) \simeq LCH^{n-i+2}_{\varepsilon}(\Lambda).$$

Here ε is the augmentation induced by L_Λ.

The definition of the augmentation induced by an exact Lagrangian filling can be found in Section 3 of [6]. Observe that Ekholm in [7] provided a fairly complete sketch of proof of Fact 4.1. In addition, Fact 4.1 will follow from the result which connects Legendrian contact homology and wrapped Floer homology for Legendrian knots in $\mathbb{R}^3$, see [11] (the proof is based on the gradient flow trees technique and the result can be extended to higher dimensions). From Fact 4.1 and the fact that Legendrian isotopy implies that there exists an exact Lagrangian cylinder, see Proposition 1.4 in [15], it follows that $\mathcal{H}^{geom}_{\Lambda}$ is a Legendrian invariant.

Note that Formula 4.3 holds for every exact Lagrangian filling L_{Λ_+} of Λ_+ obtained by gluing the positive end of an exact Lagrangian filling L_{Λ_-} to the negative end of L.

Since Λ_- is chord generic, every linearized Legendrian contact cohomology differential graded complex of Λ_- has the same (finite) number of generators. Therefore using that $\emptyset \prec^{ex}_{L_{\Lambda_-}} \Lambda_-$ and Fact 4.1 we get that there is $L^{max}_{\Lambda_-} \in \mathcal{L}(\Lambda_-)$ such that

$$\dim(H_i(L^{max}_{\Lambda_-};\mathbb{Z}_2)) \geq \dim(H_i(L_{\Lambda_-};\mathbb{Z}_2)) \tag{4.4}$$

for all $L_{\Lambda_-} \in \mathcal{L}(\Lambda_-)$. Then we construct $L^{sep}_{\Lambda_+}$ which is an exact Lagrangian cobordism obtained by gluing the positive end of $L^{max}_{\Lambda_-}$ to the negative end of L. Formulas 4.3 and 4.4 imply that

$$\dim(H_i(L^{sep}_{\Lambda_+};\mathbb{Z}_2)) > \dim(H_i(L^{max}_{\Lambda_-};\mathbb{Z}_2))$$

and hence $\mathcal{H}^{geom}_{\Lambda_-} \neq \mathcal{H}^{geom}_{\Lambda_+}$. Thus Λ_- is not Legendrian isotopic to Λ_+.

Consider $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} L$ and $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-$, where $i_1, \dots, i_k \geq i$ and i is the smallest number such that $\dim(H_i(L;\mathbb{Z}_2)) > \dim(H_i(\Lambda_-;\mathbb{Z}_2))$. Observe that $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} L$ is diffeomorphic to $L \times S^{i_1} \times \dots \times S^{i_k}$ and $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-$ is diffeomorphic to $\Lambda_- \times S^{i_1} \times \dots \times S^{i_k}$. Hence

$$\begin{aligned}(4.5)\quad H_i(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} L;\mathbb{Z}_2) &\simeq H_i(\Sigma_{S^{i_{k-1}}} \dots \Sigma_{S^{i_1}} L;\mathbb{Z}_2) \oplus H_{i-i_k,\,0}(\Sigma_{S^{i_{k-1}}} \dots \Sigma_{S^{i_1}} L;\mathbb{Z}_2)\\ &\simeq H_i(L;\mathbb{Z}_2) \oplus H_{i-i_1,\,0}(L;\mathbb{Z}_2) \oplus \dots \oplus H_{i-i_k,\,0}(\Sigma_{S^{i_{k-1}}} \dots \Sigma_{S^{i_1}} L;\mathbb{Z}_2) \simeq H_i(L;\mathbb{Z}_2) \oplus \mathbb{Z}_2^l,\end{aligned}$$

where

$$H_{i-i_j,\,0}(\,\cdot\,;\mathbb{Z}_2) \simeq \begin{cases} H_{i-i_j}(\,\cdot\,;\mathbb{Z}_2), & \text{if } i-i_j \geq 0;\\ 0, & \text{if } i-i_j < 0\end{cases}$$

and l is the number of j's such that $i_j = i$. Similarly, we see that

$$(4.6)\qquad H_i(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-;\mathbb{Z}_2) \simeq H_i(\Lambda_-;\mathbb{Z}_2) \oplus \mathbb{Z}_2^l,$$

where l is the number of j's such that $i_j = i$. Observe that here we use that Λ_- and L are connected. Formulas 4.1, 4.5 and 4.6 imply that

$$\begin{aligned}(4.7)\qquad \dim(H_i(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} L;\mathbb{Z}_2)) &= \dim(H_i(L;\mathbb{Z}_2)) + l > \dim(H_i(\Lambda_-;\mathbb{Z}_2)) + l\\ &= \dim(H_i(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-;\mathbb{Z}_2)).\end{aligned}$$

Observe that from the proof of Proposition 1.2 it follows that

$$\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_- \prec^{ex}_{\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} L} \Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_+.$$

We now take a chord generic representative $(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm)_{gen}$ in the Legendrian isotopy class of $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm$. Note that from the proof of Proposition 1.4 in [15] it follows that there are exact Lagrangian cobordisms $L^{(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm)_{gen}}_{\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm}$ from $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm$ to $(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm)_{gen}$ and $L^{\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm}_{(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm)_{gen}}$ from $(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm)_{gen}$ to $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm$ that are diffeomorphic to $\mathbb{R} \times \Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_\pm$. Hence we use the fact that $\emptyset \prec^{ex}_{\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} L_{\Lambda_-}} \Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-$, glue $L^{(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-)_{gen}}_{\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-}$ to $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} L_{\Lambda_-}$ along $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-$ and get $L_{(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-)_{gen}}$ which is an exact Lagrangian filling of $(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-)_{gen}$. In addition, if we glue $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} L$ to $L^{\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-}_{(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-)_{gen}}$ along $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-$ and to $L^{(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_+)_{gen}}_{\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_+}$ along $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_+$, then we get an exact Lagrangian cobordism $L^{(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_+)_{gen}}_{(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-)_{gen}}$ from $(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-)_{gen}$ to $(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_+)_{gen}$ which is diffeomorphic to $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} L$. From Formula 4.7 it follows that

$$\dim(H_i(L^{(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_+)_{gen}}_{(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-)_{gen}};\mathbb{Z}_2)) > \dim(H_i((\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-)_{gen};\mathbb{Z}_2)).$$

Hence the first part of Proposition 1.3 implies that $(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-)_{gen}$ is not Legendrian isotopic to $(\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_+)_{gen}$. Therefore $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_-$ is not Legendrian isotopic to $\Sigma_{S^{i_k}} \dots \Sigma_{S^{i_1}} \Lambda_+$. This finishes the proof of Proposition 1.3.

5. PROOF OF THEOREM 1.4

We first proof the following simple lemma:

Lemma 5.1. *Let Λ be a closed, orientable Legendrian submanifold of $\mathbb{R}^{2n+1}$ such that $\emptyset \prec^{ex}_{L_\Lambda} \Lambda$. Then $\Sigma_{S^m}\Lambda$ has*

1. *the topological type of $\Lambda \times S^m$,*
2. *the rotation class of $\Sigma_{S^m}\Lambda$ is determined by the rotation class of Λ, and*
3. *the Thurston-Bennequin number*

$$tb(\Sigma_{S^m}\Lambda) = \begin{cases} 2(-1)^{\frac{m}{2}} tb(\Lambda), & \text{if } m \text{ is even;} \\ 0, & \text{if } m \text{ is odd.} \end{cases}$$

Proof. We first observe that (1) and (2) are straightforward and follow from the construction of $\Sigma_{S^m}\Lambda$. Then we prove (3). Note that $L_{\Sigma_{S^m}\Lambda} := \Sigma_{S^m} L_\Lambda$ is diffeomorphic to $L_\Lambda \times S^m$. Hence we get

$$\chi(L_{\Sigma_{S^m}\Lambda}) = \chi(L_\Lambda \times S^m) = \chi(L_\Lambda)\chi(S^m) = \begin{cases} 2\chi(L_\Lambda), & \text{if } m \text{ is even;} \\ 0, & \text{if } m \text{ is odd.} \end{cases} \tag{5.1}$$

We now recall that

$$tb(\Lambda) = \begin{cases} (-1)^{\frac{n}{2}+1}\chi(L_\Lambda), & \text{if } n \text{ is even;} \\ (-1)^{\frac{(n-2)(n-1)}{2}+1}\chi(L_\Lambda), & \text{if } n \text{ is odd,} \end{cases} \tag{5.2}$$

see Remark 3.5 in [15]. Formulas 5.1 and 5.2 imply that

$$\begin{aligned} tb(\Sigma_{S^m}\Lambda) &= \begin{cases} (-1)^{\frac{m+n}{2}+1}\chi(L_{\Sigma_{S^m}\Lambda}), & \text{if } n, m \text{ are even;} \\ (-1)^{\frac{(n+m-2)(n+m-1)}{2}+1}\chi(L_{\Sigma_{S^m}\Lambda}), & \text{if } n \text{ is odd}, m \text{ is even;} \\ 0, & \text{if } m \text{ is odd;} \end{cases} \\ &= \begin{cases} 2(-1)^{\frac{m}{2}} tb(\Lambda), & \text{if } m \text{ is even;} \\ 0, & \text{if } m \text{ is odd.} \end{cases} \end{aligned} \tag{5.3}$$

□

Let T_{2k+1} be the Legendrian torus knot whose Lagrangian projection is in Figure 1 with rotation number $r(T_{2k+1}) = 0$ for $k \geq 1$. Note that T_{2k+1} admits an exact Lagrangian filling for every $k \geq 1$.

One gets an exact Lagrangian filling of T_{2k+1} by gluing the "basic" exact Lagrangian cobordisms constructed in [11], i.e., the cobordisms which correspond to Legendrian isotopy, 0-resolution at a contractible crossing in the Lagrangian projection (see Figure 2) and capping off a $tb = -1$ unknot with a disk. Observe that one can construct the cobordism which corresponds to 0-resolution at a contractible crossing using the model from Section 3.3 in [16]. In addition, the cobordism which corresponds to Legendrian isotopy has been constructed by Chantraine in [4].

Recall that $T_{2j+1} \prec^{ex}_{L^{2k+1}_{2j+1}} T_{2k+1}$ for $k > j$, see the proof of Proposition 1.6 in [15]. It is easy to see that $tb(T_{2k+1}) = 2k - 1$. We use Theorem 1.2 from [4] and get that

$$2(k-j) = tb(T_{2k+1}) - tb(T_{2j+1}) = -\chi(L^{2k+1}_{2j+1}) = 2g(L^{2k+1}_{2j+1}). \tag{5.4}$$



FIGURE 1. The knot T_{2k+1}, cf Figure 13 in [9].

Therefore L^{2k+1}_{2j+1} is a twice punctured genus $g(L^{2k+1}_{2j+1}) = k - j$ oriented surface. Hence we see that

$$\dim(H_1(L^{2k+1}_{2j+1};\mathbb{Z}_2)) = 2(k-j)+1 > 1 = \dim(H_1(T_{2j+1};\mathbb{Z}_2))$$

for $k > j$.

We then apply Propositions 1.2 and 1.3 and get that there are infinitely many pairs of exact Lagrangian cobordant and not pairwise Legendrian isotopic Legendrian $S^1 \times S^{i_1} \times \cdots \times S^{i_k}$ in $\mathbb{R}^{2(\sum_{j=1}^{k} i_j+1)+1}$. In addition, from Lemma 5.1 it follows that the classical invariants of $S^1 \times S^{i_1} \times \cdots \times S^{i_k}$'s agree if one of i_j's is odd. This finishes the proof of Theorem 1.4.

6. ALTERNATIVE APPROACH

One can use generating family cohomology (over $\mathbb{Z}_2$) instead of linearized Legendrian contact cohomology to prove a variant of Proposition 1.3 (for the basic definitions of the theory of generating families we refer to [18]):

Proposition 6.1. *Let Λ_- and Λ_+ be two closed, orientable Legendrian submanifolds of $J^1(M)$ such that $\emptyset \prec^{lag}_{L_{\Lambda_-}} \Lambda_-$, $\Lambda_- \prec^{lag}_{L} \Lambda_+$ and $\dim(H_i(L;\mathbb{Z}_2)) > \dim(H_i(\Lambda_-;\mathbb{Z}_2))$ for some i. In addition, assume that L_{Λ_-} admits a tame, compatible triple of generating families $(F_{L_{\Lambda_-}}, f^-_{\emptyset}, f^+_{\Lambda_-})$ and L is gf-compatible to Λ_-. Then Λ_- is not Legendrian isotopic to Λ_+.*

Here M is a compact manifold (or $\mathbb{R}^n$) and $J^1(M)$ is a 1-jet space of M. In addition, the property that L is gf-compatible to Λ_- means that for every tame generating family f_{Λ_-} of Λ_- there exists a tame, compatible triple of generating families $(F_L, f^-_{\Lambda_-}, f^+_{\Lambda_+})$ for L, where $f^-_{\Lambda_-}$ and f_{Λ_-} are in the same equivalence class (classes are defined up to stabilizations and fiber-preserving diffeomorphisms; for more details we refer to [18]).

One proves Proposition 6.1 by simply mimicing the proof of Proposition 1.3 and using the following observations:



FIGURE 2. The 0-resolution in the Lagrangian projection.

(i) If we glue a Lagrangian filling L_{Λ_-} of Λ_- which admits a tame, compatible triple of generating families $(F_{L_{\Lambda_-}}, f^-_{\emptyset}, f^+_{\Lambda_-})$ to L which is gf-compatible to Λ_- along Λ_-, then we get a Lagrangian filling L_{Λ_+} of Λ_+ which admits a tame, compatible triple of generating families $(F_{L_{\Lambda_+}}, f^-_{\emptyset}, f^+_{\Lambda_+})$. The way to glue two cobordisms which admit tame, compactible triples of generating families is written in [18].

(ii) Sabloff and Traynor in [18] provide a complete proof of the analoque of Fact 4.1, i.e., they prove that $GH^k(f^+_{\Lambda}) \simeq H^{k+1}(L_{\Lambda}, \Lambda; \mathbb{Z}_2)$ for a Lagrangian filling L_{Λ} of a closed, orientable Legendrian submanifold Λ with a tame, compatible triple of generating families $(F_{L_{\Lambda}}, f^-_{\emptyset}, f^+_{\Lambda})$.

(iii) $\dim(GH^i(f^+_{\Lambda_-})) \leq 2l_-$ for all i, where l_- is the number of Reeb chords of Λ_-. This follows from the description of the critical points of the difference function, see [13], [17]. Hence for a fixed i there exists a Lagrangian filling $L^{max}_{\Lambda_-}$ of Λ_- which admits a tame, compatible triple of generating families $(F_{L^{max}_{\Lambda_-}}, f^-_{\emptyset,max}, f^+_{\Lambda_-,max})$ such that

$$\begin{aligned}\dim(H_i(L^{max}_{\Lambda_-}; \mathbb{Z}_2)) = \dim(H^{n-i+1}(L^{max}_{\Lambda_-}, \Lambda_-; \mathbb{Z}_2)) = \dim(GH^{n-i}(f^+_{\Lambda_-,max})) \\ \geq \dim(GH^{n-i}(f^+_{\Lambda_-})) = \dim(H_i(L_{\Lambda_-}; \mathbb{Z}_2))\end{aligned}$$

for every Lagrangian filling L_{Λ_-} of Λ_- which admits a tame, compatible triple of generating families $(F_{L_{\Lambda_-}}, f^-_{\emptyset}, f^+_{\Lambda_-})$.

Remark 6.2. Observe that Bourgeois, Sabloff and Traynor in [3] prove that the operation of "standard" Lagrangian handle attachment can be realized as a Lagrangian cobordism which is gf-compatible to its negative end. Hence every Lagrangian cobordism obtained by gluing Lagrangian cobordisms which correspond to "standard" Lagrangian handle attachments is also gf-compatible to its negative end.

ACKNOWLEDGEMENTS

The author is deeply grateful to Frédéric Bourgeois, Baptiste Chantraine, Octav Cornea, Tobias Ekholm, Georgios Dimitroglou Rizell and Joshua Sabloff for helpful conversations and interest in his work.

REFERENCES

[1] M. Abouzaid and P. Seidel, *An open-string analogue of Viterbo functoriality*, Geometry $\&$ Topology **14** (2010), 627–718.

[2] D. Bennequin, *Entrelacements et équations de Pfaff*, Asterisque **107–108** (1983), 87–161.

[3] F. Bourgeois, J. Sabloff and L. Traynor, *Lagrangian cobordisms via generating families with applications to Legendrian geography and botany*, in preparation.

[4] B. Chantraine, *Lagrangian concordance of Legendrian knots*, Algebr. Geom. Topol. **10** (2010), 63–85.

[5] Yu. Chekanov, *Differential algebra of Legendrian links*, Invent. Math. **150** (2002), 441–483.

[6] T. Ekholm, *Rational symplectic field theory over $\mathbb{Z}_2$ for exact Lagrangian cobordisms*, J. Eur. Math. Soc. **10**(3) (2008), 641–704.

[7] T. Ekholm, *Rational SFT, linearized Legendrian contact homology, and Lagrangian Floer cohomology*, Perspectives in Analysis, Geometry, and Topology, Progress in Mathematics Volume 296, (2012), 109–145.

[8] T. Ekholm, J. Etnyre and J. Sabloff, *A duality exact sequence for Legendrian contact homology*, Duke Math J. 150 (2009), 1–75.

[9] T. Ekholm, J. Etnyre and M. Sullivan, *Non-isotopic Legendrian submanifolds in $\mathbb{R}^{2n+1}$*, J. Diff. Geom. **71** (2005), 85–128.

[10] T. Ekholm, J. Etnyre and M. Sullivan, *The contact homology of Legendrian submanifolds in $\mathbb{R}^{2n+1}$*, J. Diff. Geom. **71** (2005), 177–305.

[11] T. Ekholm, K. Honda and T. Kálmán, *Legendrian knots and exact Lagrangian cobordisms*, in preparation.

[12] Y. Eliashberg, A. Givental and H. Hofer, *Introduction to symplectic field theory*, Geom. Funct. Anal. (2000), 560–673.

[13] D. Fuchs and D. Rutherford, *Generating families and legendrian contact homology in the standard contact space*, J. Topology **4** (2011), no. 1, 190–226.

[14] K. Fukaya, P. Seidel and I. Smith, *The symplectic geometry of cotangent bundles from a categorical viewpoint*, Homological Mirror Symmetry: New Developments and Perspectives, edited by Kapustin, Kreuzer and Schlesinger, Springer Lecture Notes in Physics **757**, 2009.

[15] R. Golovko, *A note on Lagrangian cobordisms between Legendrian submanifolds of $\mathbb{R}^{2n+1}$*, preprint 2011. Available at arXiv:1108.3693.

[16] G. D. Rizell, *Legendrian ambient surgery and Legendrian contact homology*, preprint 2012. Available at arXiv:1205.5544.

[17] J. Sabloff and L. Traynor, *Obstructions to the existence and squeezing of Lagrangian cobordisms*, J. Topol. Anal. **2** (2010), no. 2, 203–232.

[18] J. Sabloff and L. Traynor, *Obstructions to Lagrangian cobordisms between Legendrian submanifolds*, preprint 2011. Available at arXiv:1109.5660.

[19] S. Tabachnikov, *An invariant of a submanifold that is transversal to a distribution*, Uspekhi Mat. Nauk **43:3**(261) (1988), 193–194.

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ LIBRE DE BRUXELLES, CP 218, BOULEVARD DU TRIOMPHE, B-1050 BRUXELLES, BELGIUM
E-mail address: `rgolovko@ulb.ac.be`
URL: `http://homepages.ulb.ac.be/~rgolovko/`